

Dynamic Kinetic Resolution-Enabled Highly Stereoselective Nucleophilic Fluoroalkylation to Access Chiral β -Fluoro Amines

Qinghe Liu, Taige Kong, Chuanfa Ni, and Jinbo Hu*



Cite This: *Org. Lett.* 2022, 24, 5982–5987



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: β -Fluorinated amine is highly desirable for biological and pharmaceutical science, because replacing a C–H bond with a C–F bond can change the physical and chemical properties of the parent molecule to a large extent but not significantly alter its overall geometry. Herein, the highly stereoselective nucleophilic monofluoromethylation of imines have been developed. It is proposed that the chelated transition state enables the chiral induction by the dynamic kinetic resolution of the chiral α -fluoro carbanions.



Amines are privileged structures present in a myriad of biologically active molecules.^{1–4} However, amino groups typically result in a poor ability to penetrate the cell membrane due to their strong basicity, thus leading to an obvious decrease in the biological activity of drug molecules.⁵ Due to the peak electronegativity of fluorine atoms, replacing a C–H bond with a C–F bond can strongly affect the basicity of nearby amino groups.^{5,6} For example, the pK_a value of the protonated 5-HT_{1D} agonist is 9.7, while that of its β -fluorinated analogue is 8.7 (Figure 1a).^{5,6} The change in the pK_a value of the drug molecule has an important impact on the kinetic metabolic properties and the affinity, which improves the cell membrane permeability, thereby improving its biological activity.^{7,8} In addition, fluorine has a considerable effect on the electron distribution of the molecule, while not significantly altering or distorting the general geometry of molecules because of the similar size of hydrogen and fluorine.⁹ Thus, ready access to β -fluoro amine is highly desirable and significant for the life sciences and pharmaceutical industry.


A number of methods for accessing β -fluoro amines have been developed, including fluoroamination of alkenes,^{10–14} nucleophilic fluorination of alcohols,^{15,16} and ring opening with metal fluorides.^{17–20} However, these methods cannot be applied for chiral β -fluoro amines, given the fact that the potential dangers of racemic drugs have been documented (Figure 1b).^{21–28} In 2007, Shibata and Toru developed an elegant catalytic enantioselective fluoromethylation with in situ-generated aldimines and FBSM, to yield chiral β -fluoro amines.²⁹ However, the reaction time was up to 2 days at -80°C , and enaminalizable ketimines were not compatible (Figure 1c-i). Hu and co-workers synthesized β -fluoro amines via diastereoselective fluoromethylation of chiral *N*-*tert*-butylsulfinyl ketimines, but with a racemic carbon center at the β -



position (due to the difficult differentiation of H/F atoms), and the yield is moderate (Figure 1c-ii).^{30,31} In addition, an enzyme-catalyzed kinetic resolution was also reported, but the drawback is the limitation of the 50% maximal yield (Figure 1c-iii).^{32,33} Our group was recently devoted to the development of nucleophilic fluoroalkylations by employing sulfoximine as the stereocontrol reagent.^{34–38} Herein, a new method for asymmetric synthesis of β -fluoro amines with a sulfoximine reagent is reported (Figure 1c-iv). Our concept relies on the dynamic kinetic resolution of α -fluoro carbanions, which delivers chiral amines with a fluorinated carbon stereogenic center at the β -position. It is worth noting that there has been no report about the reaction of imines with fluorinated sulfoximine. Previously, the reaction of ketones strictly depends on the nonchelated transition state due to the poor chelation ability of oxygen.^{36,37} However, is it possible that the strong chelation ability of nitrogen from imines can undermine the dynamic kinetic resolution process? Can we realize the dynamic kinetic resolution in a different mode? Herein, we report the first dynamic kinetic resolution-enabled highly stereoselective fluoroalkylation of imines.

To achieve high stereoselectivity, we first investigated the protecting groups of imines with (*R*)-phenyl monofluoromethyl sulfoximine 1, considering the relatively easy availability of optically pure phenyl monofluoromethyl sulfoximine (Table S1). When a phenylthio group-protected imine was used, the

Received: July 3, 2022


Published: August 8, 2022




(a) The effect of replacing the C-H with C-F on the β -position of amine(b) Representative bioactive molecules containing β -fluoro amine

(c) The state-of-the-art synthetic methods


i. Catalyst-controlled reaction:



ii. Substrate-controlled reaction:



iii. Enzyme-controlled kinetic resolution:



(d) This work

iv. Sulfoximine-controlled dynamic kinetic resolution:




Figure 1. Examples of biologically active β -fluoro amines and state-of-the-art methods for accessing chiral β -fluoro amines. Abbreviations: EGFR, bovine plasma amine oxidase; BACE1, β -site APP-cleaving enzyme 1.

reaction gave 56/44 dr (Table S1, entry 1). When we increased the steric hindrance of the protecting group (using a *tert*-butoxycarbonyl group), the reaction could give 75/25 dr (Table S1, entry 2). To our delight, using a *tert*-butylsulfonyl group (Bus)-protected imine can give 93/7 dr,³⁹ but the use of an arylsulfonyl group resulted in a decrease in the dr values (Table S1, entries 3–6).

We subsequently screened a model reaction between Bus-protected enaminalizable ketimine **4a** and sulfoximine reagent **1** for the asymmetric synthesis of β -fluoro tertiary amines (Table 1). It was found that when LiHMDS, NaHMDS, or KHMDS was used as the base, the reaction proceeded with difficulty (Table 1, entries 1–3). When the base was changed to LDA, both the yield and dr increased dramatically [42% yield, dr 98/2 (Table 1, entry 4)].⁴⁰ However, when *n*BuLi was used, a yield of 32% was afforded and two diastereoisomers were observed (dr 87/13). After the solvents had been screened, CH₂Cl₂ was found to be the optimal solvent (Table 1, entries 6–9). However, when HMPA was added, the yield and diastereoselectivity were significantly reduced (entry 10). Further optimization showed that when the **1/4a**/LDA ratio was changed to 2.5/1/2.5, we can obtain **5a** in 90% yield with 98/2 dr (Table 1, entry 13).

Then we examined the substrate scope of the reaction (Figure 2). Reactions with various aryl methyl ketimines can give **5a**–**l** in high yields (77–90%) with high diastereoselectivities (dr 95/5–98/2). Many substituents such as fluoro, chloro, bromo, iodo, methyl, isopropyl, and trifluoromethyl groups could be tolerated under the reaction conditions. The reaction was not sensitive to the position of substituents, and **5g**–**i** can be obtained in high yields with high diastereoselectivities. The reaction with a biphenyl-substituted ketimine can afford product **5m** in 74% yield with 96/4 dr. Disubstituted substrates such as electron-poor **4n** and electron-rich **4o** gave the corresponding products in 70% yield with 96/4 dr and 85% yield with 96/4 dr, respectively. **5p** was afforded in 82% yield with 98/2 dr when a naphthyl-

Table 1. Survey of the Reaction Conditions^a

entry	1/4a /base	base	solvent	yield (%) ^b	dr ^b
1	1.0/1.2/1.5	LiHMDS	CH ₂ Cl ₂	NR	ND
2	1.0/1.2/1.5	NaHMDS	CH ₂ Cl ₂	7	ND
3	1.0/1.2/1.5	KHMDS	CH ₂ Cl ₂	<5	ND
4	1.0/1.2/1.5	LDA	CH ₂ Cl ₂	42	98/2
5	1.0/1.2/1.5	<i>n</i> BuLi	CH ₂ Cl ₂	32	87/13
6	1.0/1.2/1.5	LDA	Et ₂ O	41	95/5
7	1.0/1.2/1.5	LDA	THF	35	94/6
8	1.0/1.2/1.5	LDA	DME	<5	ND
9	1.0/1.2/1.5	LDA	toluene	44	96/4
10 ^c	1.0/1.2/1.5	LDA	CH ₂ Cl ₂ /hmpa	9	50/50
11	1.0/2.0/2.5	LDA	CH ₂ Cl ₂	62	98/2
12	2.0/1.0/2.5	LDA	CH ₂ Cl ₂	90	97/3
13 ^d	2.5/1.0/2.5	LDA	CH ₂ Cl ₂	90	98/2

^aA base was added slowly to the solution of (R)-**1** in the solvent at -78 °C; 30 min later, a solution of **4a** in a solvent was added, and the mixture was stirred for an additional 3 h, followed by quenching with a saturated NH₄Cl solution. ^bThe yield and dr were determined by ¹⁹F NMR. NR, no desired reaction; ND, not determined; Bus, *tert*-butanesulfonyl. ^cAt 10/1 (v/v). ^dAt -94 °C.

substituted ketimine was used. In addition, the reactions with ketimines **4q** and **4r** afforded the corresponding products **5q** in 87% yield with 95/5 dr and **5r** in 88% yield with 96/4 dr. Moreover, a heteroaryl ketimine could also be applied to the reaction, generating **5s** in 90% yield with 97/3 dr. The reaction with an alkyl-substituted methyl ketimine proceeded smoothly, giving **5t** in 86% yield with 88/12 dr. Secondary amines could be obtained by employing aldimines, such as **5u**–**w**. The absolute configuration of **5e** was confirmed by X-ray crystal




Figure 2. Substrate scope of the highly stereoselective fluoroalkylation reaction. ^aYields refer to the isolated yield of the major diastereoisomer; dr values were determined by ¹⁹F NMR. P = Bus = *tert*-butanesulfonyl. ^bKHMDs and THF were used. The absolute configuration of **5e** was determined by single-crystal X-ray structure analysis, and those of the others were assigned by analogy.

structure analysis, and those of the other products were assigned by analogy.

To gain insights into the reaction mechanism especially in terms of the high stereoselectivity in the generation of the fluorinated carbon stereogenic center, we carried out several mechanistic experiments. First, a reaction of D₂O and the (*R*_s)-PhSO(NTBS)CHF⁻ anion was tested (Figure 3a). After




Figure 3. Mechanistic studies and proposed transition states. Bus = *tert*-butylsulfonyl. Repulsive interactions are indicated by a curved arrow.

quenching with excess aqueous ammonium chloride, 30% conversion of (*R*)-1' was observed on the basis of ¹⁹F NMR. Due to the signal overlap of the diastereomers in ¹⁹F spectra, we conducted ¹H NMR analysis and found that the diastereomeric ratio was 51/49, which indicated that carbanions in both *R* and *S* configurations at the fluorocarbon center were formed without selectivity (part 5 of the Supporting Information). Second, we tried to examine whether the nucleophilic reaction is reversible. Upon treatment of (*R*)-1 with *n*BuLi and 4a, compound 5a' was achieved with three diastereomers (Figure 3b, 85/8/7 dr). Subsequently, compound 5a' was recovered fully without any changes in dr after being treated with LDA at -94 °C for 0.5 h (Figure 3c), which implied that this reaction was not reversible. We therefore proposed the high stereoselectivity at the fluorinated stereocenter was caused by the dynamic kinetic resolution of the participating carbanions (Figure 1d). It is obvious that the facial selectivity of the addition can be attributed to the chiral induction from the sulfur stereogenic center to the fluorine-bearing carbon stereogenic center.

By comparing different chair- or boat-like chelating transition states A–C, we rationalized the stereocontrol mode of the reaction (Figure 4d). First, because of the flagpole interaction of oxygen and the Ar group, boat-like transition state B is disfavored.^{41a} Second, due to the 1,3-diaxial interaction of Ar and the phenyl group, chair-like chelating transition state C is energetically unfavorable,^{41b} although it leads to the experimentally observed major diastereomer 5a (Figure 4d, C). Our proposed transition state A can be supported by the result that the addition of HMPA, which prevents the chelation of the sulfoximine oxygen atom with the metal (for example, lithium cation), can significantly reduce the yield and diastereoselectivity. It is




Figure 4. Extension of the reaction and product.

parallel to what is proposed by Pyne regarding the reactions of ketones and sulfoximine⁴² (Table 1, entry 10).

On this basis, the highly nucleophilic monofluoroalkylation can also be extended to the difluoroalkylation of imines with high dr. The results are shown in Figure 4a. It was found that 1-(naphthalen-2-yl)ethyl imine 4p reacted with the (S)-phenyl difluoromethyl sulfoximine reagent to afford addition product 8a in 88% yield with 99/1 dr. The reaction tolerated 1-(benzo[d][1,3]dioxol-5-yl)ethyl, 1-phenylpropyl, and 1-(2-methoxyphenyl)ethyl imines, to generate 8b–d, respectively, in high yields and diastereoselectivities, and the configuration of 8a was confirmed by X-ray crystal analysis. To obtain the monofluoromethylated free amine, 5a could undergo deprotection of the Bus group with AlCl_3 and reductive C–S bond cleavage with aluminum amalgam⁴³ to yield free amine 9a in 63% yield and 98/2 er (Figure 4b). Free amines can allow different modifications and provide the possibility of accessing chiral amine derivatives, especially those bioactive molecules.^{21–28}

In conclusion, we report the first dynamic kinetic resolution-enabled stereoselective fluoroalkylation of imines with a fluoromethyl sulfoximine reagent 1. The highly stereoselective nucleophilic fluoroalkylation of imines with a broad substrate scope was developed. Mechanistic studies showed that the chiral induction from the sulfur stereogenic center of the sulfoximine to the fluorine-bearing carbon stereogenic center was enabled by a dynamic kinetic resolution of the α -fluorinated carbanion. The reaction can also be extended to the difluoroalkylation reaction, which further expands the scope of this protocol.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.2c02250>.

Detailed experimental procedures and spectra data for all new compounds (PDF)

Accession Codes

CCDC 2166836–2166837 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The

Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

■ AUTHOR INFORMATION

Corresponding Author

Jinbo Hu – Key Laboratory of Organofluorine Chemistry, Center for Excellence in Molecular Synthesis, Shanghai Institute of Organic Chemistry, University of Chinese Academy of Sciences, Chinese Academy of Sciences, Shanghai 200032, P. R. China; orcid.org/0000-0003-3537-0207; Email: jinbohu@sioc.ac.cn

Authors

Qinghe Liu – Key Laboratory of Organofluorine Chemistry, Center for Excellence in Molecular Synthesis, Shanghai Institute of Organic Chemistry, University of Chinese Academy of Sciences, Chinese Academy of Sciences, Shanghai 200032, P. R. China; orcid.org/0000-0003-3125-7135

Taige Kong – Key Laboratory of Organofluorine Chemistry, Center for Excellence in Molecular Synthesis, Shanghai Institute of Organic Chemistry, University of Chinese Academy of Sciences, Chinese Academy of Sciences, Shanghai 200032, P. R. China

Chuanfa Ni – Key Laboratory of Organofluorine Chemistry, Center for Excellence in Molecular Synthesis, Shanghai Institute of Organic Chemistry, University of Chinese Academy of Sciences, Chinese Academy of Sciences, Shanghai 200032, P. R. China

Complete contact information is available at:

<https://pubs.acs.org/doi/10.1021/acs.orglett.2c02250>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Financial support was provided by the National Key Research and Development Program of China (2021YFF0701700), the National Natural Science Foundation of China (21632009), the Key Programs of the Chinese Academy of Sciences (KGZD-EW-T08), and the Key Research Program of Frontier Sciences of CAS (QYZDJ-SSW-SLH049).

■ REFERENCES

- (1) Kobayashi, S.; Ishitani, H. Catalytic Enantioselective Addition to Imines. *Chem. Rev.* **1999**, *99*, 1069–1094.
- (2) Ellman, J. A.; Owens, T. D.; Tang, T. P. N-tert-Butanesulfinyl Imines: Versatile Intermediates for the Asymmetric Synthesis of Amines. *Acc. Chem. Res.* **2002**, *35*, 984–995.
- (3) Xie, J.-H.; Zhu, S.-F.; Zhou, Q.-L. Recent advances in transition metal-catalyzed enantioselective hydrogenation of unprotected enamines. *Chem. Soc. Rev.* **2012**, *41*, 4126–4139.
- (4) Breuer, M.; Ditrach, K.; Habicher, T.; Hauer, B.; Keßler, M.; Stürmer, R.; Zelinski, T. Industrial Methods for the Production of Optically Active Intermediates. *Angew. Chem., Int. Ed.* **2004**, *43*, 788–824.
- (5) Böhm, H.-J.; Banner, D.; Bendels, S.; Kansy, M.; Kuhn, B.; Müller, K.; Obst-Sander, U.; Stahl, M. Fluorine in Medicinal Chemistry. *ChemBioChem.* **2004**, *5*, 637–643.
- (6) *MedChem Database*, Biobyte Corp. and Pomona College, 2002.
- (7) Lankin, D. C.; Chandrakumar, N. S.; Rao, S. N.; Spangler, D. P.; Snyder, J. P. Protonated 3-fluoropiperidines: an unusual fluoro directing effect and a test for quantitative theories of solvation. *J. Am. Chem. Soc.* **1993**, *115*, 3356–3357.

- (8) van Niel, M. B.; Collins, I.; Beer, M. S.; Broughton, H. B.; Cheng, S. K. F.; Goodacre, S. C.; Heald, A.; Locker, K. L.; MacLeod, A. M.; Morrison, D.; Moyes, C. R.; O'Connor, D.; Pike, A.; Rowley, M.; Russell, M. G. N.; Sohal, B.; Stanton, J. A.; Thomas, S.; Verrier, H.; Watt, A. P.; Castro, J. L. Fluorination of 3-(3-(Piperidin-1-yl)propyl)indoles and 3-(3-(Piperazin-1-yl)propyl)indoles Gives Selective Human 5-HT1D Receptor Ligands with Improved Pharmacokinetic Profiles. *J. Med. Chem.* **1999**, *42*, 2087–2104.
- (9) (a) O'Hagan, D. Understanding organofluorine chemistry. An introduction to the C-F bond. *Chem. Soc. Rev.* **2008**, *37*, 308–319. (b) Qing, F.-L.; Liu, X.-Y.; Ma, J.-A.; Shen, Q.; Song, Q.; Tang, P. A Fruitful Decade of Organofluorine Chemistry: New Reagents and Reactions. *CCS Chem.* **2022**, DOI: 10.31635/ccschem.022.202201935. (c) Chen, N.; Lei, J.; Wang, Z.; Liu, Y.; Sun, K.; Tang, S. Construction of Fluoro-containing Heterocycles Mediated by Free Radicals. *Chin. J. Org. Chem.* **2022**, *42*, 1061–1084.
- (10) Zhang, Q.; Zheng, G.; Zhang, Q.; Li, Y.; Zhang, Q. Metal-Free Three-Component Regioselective Aminofluorination of Styrene Derivatives. *J. Org. Chem.* **2017**, *82*, 8258–8266.
- (11) Qiu, S.; Xu, T.; Zhou, J.; Guo, Y.; Liu, G. Palladium-Catalyzed Intermolecular Aminofluorination of Styrenes. *J. Am. Chem. Soc.* **2010**, *132*, 2856–2857.
- (12) Xiong, T.; Zhang, Q. New amination strategies based on nitrogen-centered radical chemistry. *Chem. Soc. Rev.* **2016**, *45*, 3069–3087.
- (13) Zhang, H.; Song, Y.; Zhao, J.; Zhang, J.; Zhang, Q. Regioselective Radical Aminofluorination of Styrenes. *Angew. Chem., Int. Ed.* **2014**, *53*, 11079–11083.
- (14) Yadav, J. S.; Subba Reddy, B. V.; Narasimha Chary, D.; Chandrakanth, D. InX3-catalyzed haloamidation of vinyl arenes: a facile synthesis of α -bromo- and α -fluoroamides. *Tetrahedron Lett.* **2009**, *50*, 1136–1138.
- (15) Ducep, J. B.; Heintzelmann, B.; Jund, K.; Lesur, B.; Schleimer, M.; Zimmermann, P. R. Synthesis of (2S,5S)-5-fluoromethylornithine; a potent inhibitor of ornithine aminotransferase. *Tetrahedron: Asym.* **1997**, *8*, 327–335.
- (16) Schumacher, D. P.; Clark, J. E.; Murphy, B. L.; Fischer, P. A. An efficient synthesis of florfenicol. *J. Org. Chem.* **1990**, *55*, S291–S294.
- (17) Posakony, J. J.; Tewson, T. J. Substitution Reactions of Hindered Cyclic Sulfamides. *Synthesis* **2002**, *2002*, 859–864.
- (18) Fan, R.-H.; Zhou, Y.-G.; Zhang, W.-X.; Hou, X.-L.; Dai, L.-X. Facile Preparation of β -Fluoro Amines by the Reaction of Aziridines with Potassium Fluoride Dihydrate in the Presence of Bu₄NHSO₄. *J. Org. Chem.* **2004**, *69*, 335–338.
- (19) Alvernhe, G. M.; Ennakoua, C. M.; Lacombe, S. M.; Laurent, A. J. Ring opening of aziridines by different fluorinating reagents: three synthetic routes to α,β -fluoro amines with different stereochemical pathways. *J. Org. Chem.* **1981**, *46*, 4938–4948.
- (20) Adler, P.; Teskey, C. J.; Kaiser, D.; Holy, M.; Sitte, H. H.; Maulide, N. α -Fluorination of carbonyls with nucleophilic fluorine. *Nat. Chem.* **2019**, *11*, 329–334.
- (21) Kollonitsch, J.; Patchett, A. A.; Marburg, S.; Maycock, A. L.; Perkins, L. M.; Doldouras, G. A.; Duggan, D. E.; Aster, S. D. Selective inhibitors of biosynthesis of aminergic neurotransmitters. *Nature* **1978**, *274*, 906–908.
- (22) Grunewald, G. L.; Caldwell, T. M.; Li, Q.; Slavica, M.; Criscione, K. R.; Borchardt, R. T.; Wang, W. Synthesis and Biochemical Evaluation of 3-Fluoromethyl-1,2,3,4-tetrahydroisoquinolines as Selective Inhibitors of Phenylethanolamine N-Methyltransferase versus the α 2-Adrenoceptor. *J. Med. Chem.* **1999**, *42*, 3588–3601.
- (23) Silverman, R. B.; Nanavati, S. M. Selective inhibition of gamma-aminobutyric acid aminotransferase by (3R,4R), (3S,4S)- and (3R,4S), (3S,4R)-4-amino-5-fluoro-3-phenylpentanoic acids. *J. Med. Chem.* **1990**, *33*, 931–936.
- (24) Bey, P.; Gerhart, F.; Van Dorsselaer, V.; Danzin, C. α -(Fluoromethyl)dehydroornithine and α -(fluoromethyl)-dehydroputrescine analogs as irreversible inhibitors of ornithine decarboxylase. *J. Med. Chem.* **1983**, *26*, 1551–1556.
- (25) Krane Thvedt, T. H.; Kaasa, K.; Sundby, E.; Charnock, C.; Hoff, B. H. Chiral N-benzyl-N-methyl-1-(naphthalen-1-yl)-ethanamines and their in vitro antifungal activity against *Cryptococcus neoformans*, *Trichophyton mentagrophytes* and *Trichophyton rubrum*. *Eur. J. Med. Chem.* **2013**, *68*, 482–496.
- (26) Bugge, S.; Kaspersen, S. J.; Larsen, S.; Nonstad, U.; Bjørkøy, G.; Sundby, E.; Hoff, B. H. Structure-activity study leading to identification of a highly active thienopyrimidine based EGFR inhibitor. *Eur. J. Med. Chem.* **2014**, *75*, 354–374.
- (27) Hilpert, H.; Guba, W.; Woltering, T. J.; Wostl, W.; Pinard, E.; Mauser, H.; Mayweg, A. V.; Rogers-Evans, M.; Humm, R.; Krummenacher, D.; Muser, T.; Schnider, C.; Jacobsen, H.; Ozmen, L.; Bergadano, A.; Banner, D. W.; Hochstrasser, R.; Kuglstatter, A.; David-Pierson, P.; Fischer, H.; Polara, A.; Narquizian, R. β -Secretase (BACE1) Inhibitors with High in Vivo Efficacy Suitable for Clinical Evaluation in Alzheimer's Disease. *J. Med. Chem.* **2013**, *56*, 3980–3995.
- (28) Peddie, V.; Pietsch, M.; Bromfield, K. M.; Pike, R. N.; Duggan, P. J.; Abell, A. D. Fluorinated β^2 - and β^3 -Amino Acids: Synthesis and Inhibition of α -Chymotrypsin. *Synthesis* **2010**, *2010*, 1845–1859.
- (29) Mizuta, S.; Shibata, N.; Goto, Y.; Furukawa, T.; Nakamura, S.; Toru, T. Cinchona Alkaloid-Catalyzed Enantioselective Monofluoromethylation Reaction Based on Fluorobis(phenylsulfonyl)methane Chemistry Combined with a Mannich-type Reaction. *J. Am. Chem. Soc.* **2007**, *129*, 6394–6395.
- (30) Liu, J.; Zhang, L.; Hu, J. Stereoselective Monofluoromethylation of N-tert-Butylsulfinyl Ketimines Using Pregenerated Fluoro(phenylsulfonyl)methyl Anion. *Org. Lett.* **2008**, *10*, 5377–5380.
- (31) Li, Y.; Ni, C.; Liu, J.; Zhang, L.; Zheng, J.; Zhu, L.; Hu, J. Stereoselective Nucleophilic Monofluoromethylation of N-(tert-Butanesulfinyl)imines with Fluoromethyl Phenyl Sulfone. *Org. Lett.* **2006**, *8*, 1693–1696.
- (32) Cuetos, A.; García-Ramos, M.; Fischereder, E.-M.; Díaz-Rodríguez, A.; Grogan, G.; Gotor, V.; Kroutil, W.; Lavandera, I. Catalytic Promiscuity of Transaminases: Preparation of Enantioenriched β -Fluoroamines by Formal Tandem Hydrodefluorination/Deamination. *Angew. Chem., Int. Ed.* **2016**, *55*, 3144–3147.
- (33) Thvedt, T. H. K.; Fuglseth, E.; Sundby, E.; Hoff, B. H. Enantioenriched 1-aryl-2-fluoroethylamines. Efficient lipase-catalysed resolution and limitations to the Mitsunobu inversion protocol. *Tetrahedron* **2010**, *66*, 6733–6743.
- (34) Shen, X.; Hu, J. Fluorinated Sulfoximines: Preparation, Reactions and Applications. *Eur. J. Org. Chem.* **2014**, *2014*, 4437–4451.
- (35) Liu, Q.; Ni, C.; Wang, Q.; Meng, D.; Hu, J. Difluoromethyl 2-Pyridyl Sulfoximine: A Stereoselective Nucleophilic Reagent for Difluoro(aminosulfinyl)methylation and Difluoro(aminosulfonyl)-methylation. *CCS Chem.* **2022**, DOI: 10.31635/ccschem.022.202101634.
- (36) Liu, Q.; Shen, X.; Ni, C.; Hu, J. Stereoselective Carbonyl Olefination with Fluorosulfoximines: Facile Access to Z or E Terminal Monofluoroalkenes. *Angew. Chem., Int. Ed.* **2017**, *56*, 619–623.
- (37) Shen, X.; Miao, W.; Ni, C.; Hu, J. Stereoselective Nucleophilic Fluoromethylation of Aryl Ketones: Dynamic Kinetic Resolution of Chiral α -Fluoro Carbanions. *Angew. Chem., Int. Ed.* **2014**, *53*, 775–779.
- (38) Shen, X.; Liu, Q.; Luo, T.; Hu, J. Nucleophilic Difluoromethylation of Epoxides with PhSO(NTBS)CF₂H by a Preorganization Strategy. *Chem. - Eur. J.* **2014**, *20*, 6795–6800.
- (39) Ortiz, P.; Collados, J. F.; Jumde, R. P.; Otten, E.; Harutyunyan, S. R. Copper-Catalyzed Enantioselective Alkylation of Enolizable Ketimines with Organomagnesium Reagents. *Angew. Chem., Int. Ed.* **2017**, *56*, 3041–3044.
- (40) Reich, H. J. Role of Organolithium Aggregates and Mixed Aggregates in Organolithium Mechanisms. *Chem. Rev.* **2013**, *113*, 7130–7178.
- (41) (a) Shen, X.; Zhang, W.; Ni, C.; Gu, Y.; Hu, J. Tuning the Reactivity of Difluoromethyl Sulfoximines from Electrophilic to Nucleophilic: Stereoselective Nucleophilic Difluoromethylation of

Aryl Ketones. *J. Am. Chem. Soc.* **2012**, *134*, 16999–17002.
(b) Hwang, K. J.; Logusch, E. W.; Brannigan, L. H.; Thompson, M. R. Diastereoselective 1,2-addition of lithiated N-(trialkylsilyl)-methylphenylsulfoximines to aldehydes: preparation of 1,4,3-oxathiazin-2(6H)-one 4-oxides as novel cyclic sulfoximines. *J. Org. Chem.* **1987**, *52*, 3435–3441.

(42) Pyne, S. G.; Dong, Z.; Skelton, B. W.; White, A. H. Chiral sulfur compounds. Part 25. Diastereoselective 1,2-additions of lithiated (+)-(S)-N-tert-butyl-diphenylsilyl-S-methyl-S-phenylsulfoximine to ketones. *J. Chem. Soc., Perkin Trans. 1* **1994**, 2607–2613.

(43) Li, B.; Chen, J.; Zhang, Z.; Gridnev, I. D.; Zhang, W. Nickel-Catalyzed Asymmetric Hydrogenation of N-Sulfonyl Imines. *Angew. Chem., Int. Ed.* **2019**, *58*, 7329–7334.

Recommended by ACS

N-Iodosuccinimide-Mediated Synthesis of Benzo-Fused Bisimidazoles Enabled by a One-Pot Tandem Reaction of Fluorinated Propargyl Amidines

Shan Li, Yajun Li, *et al.*

MARCH 16, 2023

THE JOURNAL OF ORGANIC CHEMISTRY

READ 

Dual Role of (NH₄)₂CO₃ Enables Defluorinative Synthesis of β -Fluoroalkylated Aminovinyl Ketones

Yu-Lan Chen, Xue-Qiang Chu, *et al.*

DECEMBER 02, 2022

ORGANIC LETTERS

READ 

Catalytic Enantioconvergent Conjugate Addition of Organosilanes via a Strategy of Fluorodesilylation

Jisheng Luo, Li Deng, *et al.*

DECEMBER 13, 2022

JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

Gram-Scale Asymmetric Synthesis of Fluorinated Amino Acids Using a Chiral Nickel(II) Complex

Thomas Hohmann, Beate Koksche, *et al.*

AUGUST 04, 2022

THE JOURNAL OF ORGANIC CHEMISTRY

READ 

Get More Suggestions >